



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

DETERMINATION OF THE ABSTRACT GROUPS OF ORDER p^2qr ;

p, q, r BEING DISTINCT PRIMES*

OLIVER E. GLENN

Since the publication† in 1899 of Professor MILLER's "Report on recent progress in the theory of groups of finite order," WESTERN‡ has published his determination of the groups of order p^3q , and LE VASSEUR§ has discussed the order p^2q^2 . This paper is devoted to the determination of all groups of the order p^2qr . It thus completes the discussion of the problem of groups whose orders are products of four primes. ||

With the exception of the group of order $2^2 \cdot 3 \cdot 5$, simply isomorphic with the icosahedron-group, all groups of order p^2qr are solvable. The maximal self-conjugate subgroups will therefore serve as the basis of classification. The twelve possible arrangements of the factors of composition are

- (1) $ppqr$, (2) $pprq$, (3) $pqpr$, (4) $pqrp$, (5) $prpq$, (6) $prqp$,
(7) $qppr$, (8) $qprr$, (9) $qrpp$, (10) $rqpp$, (11) $rppq$, (12) $rpqp$.

If for a given type of group precisely the arrangements $(i), (j), (k), \dots$, of the factors of composition are possible, then we symbolize ¶ the group (i, j, k, \dots) . Two groups having distinct symbols cannot be simply isomorphic.

The group G always contains a maximal invariant subgroup** of order p^2q , and may contain maximal subgroups†† of order p^2r and pqr . We shall discuss

* Presented to the American Mathematical Society (New York) February 25, 1905. Received for publication July 1, 1905.

† Bulletin, American Mathematical Society, vol. 1 (1899), p. 227.

‡ Proceedings of the London Mathematical Society, vol. 30 (1899), p. 209.

§ Annales Toulouse, 1903, p. 63. Comptes Rendus, vol. 128 (1899), p. 1152, and lithographed book.

|| HÖLDER, Mathematische Annalen, vol. 43 (1893), p. 335. BURNSIDE, *Finite Groups*, p. 81. HÖLDER, Göttinger Nachrichten (1895), p. 211.

¶ Additional abbreviations used throughout are the following: P, Q, \dots , operations of order p, q, \dots ; $H_{h,i}$, a maximal invariant subgroup of G , order h and type i ; $\rho_{\Omega,h}$, number of subgroups of G , order h , permutable with Ω ; N_h , number of subgroups of G of order h .

** FROBENIUS, Berliner Sitzungsberichte, vol. 1 (1895), p. 170.

†† HÖLDER, loc. cit., COLE and GLOVER, American Journal of Mathematics, vol. 15 (1893), p. 202. BURNSIDE, *Theory of Groups*, p. 63.

in detail in this paper only two classes of groups: those possessing invariant subgroups of both the types H_{p^2q} and H_{p^2r} , and those possessing maximal invariant subgroups of the type H_{p^2q} only. A detailed summary of the results obtained in the other classes is given at the end. We shall thus be concerned principally with the subgroups $H_{p^2\sigma}$ ($\sigma = q, r$) all types of which are given in the following table, in which τ denotes the number of distinct types, while (p) signifies (modulo p):

$H_{p^2\sigma, i}$	$S_3^{-1}SS_3$	$S_3^{-1}S_1S_3$	$S_3^{-1}S_2S_3$	$S_2^{-1}S_1S_2$	Parameters	τ
$i = I$	S	.	.	.		1
II	.	S_1	S_2	S_1		1
III	.	S_1^a	S_2	S_1	$\alpha^\sigma \equiv 1(p), p \equiv 1(\sigma)$	1
IV	S^a	.	.	.	$\alpha^\sigma \equiv 1(p^2), p \equiv 1(\sigma)$	1
V	.	S_1^a	$S_2^{a^h}$	S_1	$\alpha^\sigma \equiv 1(p), p \equiv 1(\sigma) \text{ 1 or } \frac{1}{2}[\sigma+1]$	
VI	.	S_2	$S_1^{-1}S_2^{i^p+i}$	S_1	$* i^\sigma \equiv 1(p), p \equiv -1(\sigma)$	1

$\sigma = q, r; S^{p^2} = 1, S_1^p = S_2^p = 1, S_3^\sigma = 1.$

§ 1. Determination of $\rho_{\Omega, h}$.

By SYLOW's theorem,† $N_\sigma = qr/\sigma, p, p^2, pqr/\sigma, p^2qr/\sigma$ or 1. If $N_\sigma = 1$ then $\rho_{\Omega, \sigma} = 1$, Ω being any operator of prime order in G . When $N_\sigma > 1$, the result of transforming the single conjugate set of N_σ subgroups

$$g_1, g_2, g_3, \dots, g_{N_\sigma}$$

by Ω is to permute them among themselves. Hence

$$\Omega^{-1}(g_1, g_2, \dots, g_{N_\sigma})\Omega = \begin{pmatrix} g_1 & g_2 & \dots & g_{N_\sigma} \\ g_{i_1} & g_{i_2} & \dots & g_{i_{N_\sigma}} \end{pmatrix} = J_{\Omega, \sigma}.$$

It follows that $J_{\Omega, \sigma}^\omega = 1$ and

$$(1) \quad N_\sigma - \rho_{\Omega, \sigma} \equiv 0 \pmod{\omega}; \quad \rho_{\Omega, \sigma} \geq 1.$$

Next let $\omega = \sigma$. Then $N_p = (p^2 - 1)/(p - 1) = p + 1$, and

$$(2) \quad p + 1 - \rho_{\Omega, p} \equiv 0 \pmod{\sigma}.$$

Hence either $\rho_{\Omega, p} = 0$ or else $\rho_{\Omega, p} \geq 2$ ($\omega = q, r$). Now if the subgroup I_{p^2} of $H_{p^2\sigma, i}$ is cyclical the order of its group of isomorphisms is

$$I = \phi(p^2) = p(p - 1).$$

* Throughout the paper i denotes a non-integral mark of the $GF[p^2]$. Thus $i^\sigma \equiv 1(p)$ is an abbreviation for $i^\sigma \equiv 1 \pmod{p, P}$, P being any quadratic function irreducible modulo p .

† SYLOW, *Mathematische Annalen*, vol. 5 (1872).

If I_{p^2} is of type $[1, 1]$ its group of isomorphisms is simply isomorphic with the congruence group $\{S_1, S_2 \dots\}$ of order $I = p(p-1)^2(p+1)$, where S_1 is

$$y_1 \equiv a_{11}x_1 + a_{12}x_2, \quad y_2 \equiv a_{21}x_1 + a_{22}x_2 \pmod{p},$$

or say

$$S_1 = (a_{11}x_1 + a_{12}x_2, a_{21}x_1 + a_{22}x_2).$$

Since Ω corresponds to an isomorphism of G , $\{\Omega\}$ corresponds to a subgroup of the group of isomorphisms of G and ω divides I . Hence when I_{p^2} is cyclical, or when $I_{p^2} = [1, 1]$ and $p \equiv 1(\sigma)$, $\rho_{\Omega, p} \geq 2$. But when $p \equiv -1(\sigma)$ and p is odd, $\rho_{\Omega, p} = 0$. Also since $\rho_{\Omega, \sigma} \geq 1$, $J_{Q, \sigma}$ and $J_{R, \sigma}$ may be permutable. If

$$S_2 = (b_{11}x_1 + b_{12}x_2, b_{21}x_1 + b_{22}x_2)$$

the necessary and sufficient conditions that $S_1 S_2 = S_2 S_1$ are

$$(3) \quad \delta_{12} = \begin{vmatrix} a_{12} & b_{12} \\ a_{11} - a_{22} & b_{11} - b_{22} \end{vmatrix} \equiv 0,^* \quad \delta'_{12} = \begin{vmatrix} a_{21} & b_{21} \\ a_{11} - a_{22} & b_{11} - b_{22} \end{vmatrix} \equiv 0,$$

$$d_{12} = \begin{vmatrix} a_{12} & a_{21} \\ b_{12} & b_{21} \end{vmatrix} \equiv 0.$$

§ 2. Class (9, 10), $p > q > r$.

We now consider the groups whose symbol is (9, 10), having the maximal subgroups $H_{p^2q, i}$ and $H_{p^2r, j}$ ($i, j = \text{IV, V, VI}$). Since I_{p^2} is invariant in G the existence of a subgroup of type IV excludes the possibility of a subgroup of type V or VI, and vice versa. There are thus five cases to consider.

[1] $i = j = \text{IV}$. Here $I_{p^2} = \{P\}$ is cyclical and P may be regarded as the generator of order p^2 in both H -sub-groups. Since $\rho_{Q, r} \geq 1$, we may choose $\{R\}$ permutable with Q and, since $q > r$, $QR = RQ$, so that G is defined by

$$P^{p^2} = Q^q = R^r = 1, \quad Q^{-1}PQ = P^a, \quad R^{-1}PR = P^\beta, \quad QR = RQ;$$

or for brevity $G = (\alpha : \beta : 1)$, where

$$\alpha^q \equiv 1, \quad \beta^r \equiv 1(p^2), \quad p \equiv 1(qr), \quad \tau = 1.$$

[2] $i = j = \text{V}$. Let $H_{p^2q, i} = \{P'_1, P'_2, Q\}$, $H_{p^2r, j} = \{P_1, P_2, R\}$, wherein $QR = RQ$. We may write

$$R^{-1}P_1R = P_1^a, \quad R^{-1}P_2R = P_2^\beta, \quad \alpha^r \equiv 1(p), \quad \beta \equiv \alpha^h.$$

$$Q^{-1}P_1Q = P_1^{a_{11}}P_2^{a_{21}}, \quad Q^{-1}P_2Q = P_1^{a_{12}}P_2^{a_{22}},$$

and from the permutable isomorphisms of I_{p^2}

$$J_Q = \begin{pmatrix} P_1^{x_1} P_2^{x_2} \\ P_1^{a_{11}x_1 + a_{12}x_2} P_2^{a_{21}x_1 + a_{22}x_2} \end{pmatrix}, \quad J_R = \begin{pmatrix} P_1^{x_1} P_2^{x_2} \\ P_1^{ax_1} P_2^{\beta x_2} \end{pmatrix},$$

* All congruences are taken modulo p unless otherwise indicated

$$(4) \quad \delta_{12} = a_{12}(\alpha - \beta) \equiv 0, \quad \delta'_{12} = a_{21}(\alpha - \beta) \equiv 0.$$

Reserving for later treatment the ambiguous case $h = 1$, we deduce $a_{12} \equiv a_{21} \equiv 0$. Suppose next that

$$R^{-1}P'_i R = P_1^{b_{1i}} P_2^{b_{2i}} \quad (i = 1, 2).$$

Then

$$(5) \quad \begin{aligned} (RQ)^{-1}P'_1(RQ) &= P_1^{a_{11}b_{11}} P_2^{a_{22}b_{21}} = (QR)^{-1}P'_1(QR) = P_1^{\gamma b_{11}} P_2^{\gamma b_{21}}, \\ b_{11}(a_{11} - \gamma) &\equiv 0, \quad b_{21}(a_{22} - \gamma) \equiv 0, \quad \gamma^q \equiv 1, \\ b_{12}(a_{11} - \delta) &\equiv 0, \quad b_{22}(a_{22} - \delta) \equiv 0, \quad \delta \equiv \gamma^k. \end{aligned}$$

Thus when $h \neq 1$, $k \neq 1$ we have one of the two equivalent results

$$a_{11} \equiv \gamma, \quad a_{22} \equiv \delta \quad \text{or} \quad a_{11} \equiv \delta, \quad a_{22} \equiv \gamma.$$

In case $h \neq 1$, $k = 1$, the set (5) becomes

$$\begin{aligned} b_{11}(a_{11} - \gamma) &\equiv 0, \quad b_{21}(a_{22} - \gamma) \equiv 0, \\ b_{12}(a_{11} - \gamma) &\equiv 0, \quad b_{22}(a_{22} - \gamma) \equiv 0, \end{aligned}$$

and there are three possibilities to consider, viz.,

- (i) $a_{11} \neq \gamma, \quad b_{11} \equiv 0, \quad b_{12} \equiv 0, \quad b_{21} \neq 0, \quad b_{22} \neq 0, \quad a_{22} \equiv \gamma;$
- (ii) $a_{11} \equiv \gamma, \quad a_{22} \neq \gamma, \quad b_{21} \equiv b_{22} \equiv 0, \quad b_{11} \neq 0, \quad b_{12} \neq 0;$
- (iii) $a_{11} \equiv \gamma, \quad a_{22} \equiv \gamma.$

Case (i) implies

$$\begin{aligned} R^{-1}P'_1 R &= P_2^{b_{21}}, \quad R^{-1}P'_2 R = P_2^{b_{22}}, \\ R^{-1}P_1^{b_{22}} R &= R^{-1}P_2^{b_{21}} R \quad \text{or} \quad P_1^{b_{22}} = P_2^{b_{21}}, \end{aligned}$$

contrary to the independence of P'_1 and P'_2 . Likewise, case (ii) is excluded. Hence $a_{11} \equiv a_{22} \equiv \gamma$.

In a similar manner, when $h = 1$, $k \neq 1$, we get $a_{11} \equiv a_{22} \equiv \alpha$.

Next let $h = 1$, $k = 1$, so that

$$R^{-1}P_i R = P_i^\alpha, \quad Q^{-1}P_i Q = P_i^\gamma \quad (i = 1, 2).$$

One of the operations P'_1, P'_2 must be independent of P_1 . As $\gamma^q \equiv 1 \pmod{p}$, we may assume that P_1 and P'_2 are independent. These will generate I_{p^2} , so that

$$Q^{-1}P_1 Q = P_1^{a_{11}} P_2^{a_{21}}, \quad R^{-1}P'_2 R = P_1^{b_{12}} P_2^{b_{22}}.$$

The abelian conditions from J_Q and J_R are [Eq. (3)]

$$\delta_{12} = b_{12}(a_{11} - \delta) \equiv 0, \quad \delta'_{12} = a_{21}(b_{22} - \alpha) \equiv 0, \quad d_{12} = a_{21}b_{12} \equiv 0.$$

Thus three possibilities arise, viz.,

$$(i) \quad a_{21} \equiv 0, \quad b_{12} \equiv 0, \quad a_{11} \equiv \delta;$$

$$(ii) \quad a_{21} \not\equiv 0, \quad b_{12} \equiv 0, \quad b_{22} \equiv \alpha;$$

$$(iii) \quad a_{21} \equiv 0, \quad b_{12} \equiv 0.$$

For (i), let $P'_1 = P_1^x P_2'^y$, $P_2 = P_1^z P_2'^w$, whence

$$Q^{-1}P'_1Q = P_1^{rx}P_2'^{y\gamma} = P_1^{\delta x}P_2'^{\delta y},$$

$$R^{-1}P_2R = P_1^{z\beta}P_2'^{w\beta} = P_1^{\alpha z + b_{12}w}P_2'^{b_{22}w},$$

$$(\gamma - \delta)x \equiv 0, \quad (\gamma - \delta)y \equiv 0,$$

$$w(b_{22} - \beta) \equiv 0, \quad z(\alpha - \beta) + b_{12}w \equiv 0.$$

Hence $\gamma \equiv \delta$ and $k = 1$; but as P_1, P_2 are independent, $w \not\equiv 0$, $b_{22} \equiv \beta$, $\alpha \not\equiv \beta$ and $h \not\equiv 1$, contrary to hypothesis. Since (ii) is likewise excluded, we have $a_{21} \equiv b_{12} \equiv 0$,

$$Q^{-1}P_1Q = P_1^{a_{11}}, \quad R^{-1}P'_2R = P_2'^{b_{22}},$$

$$x(a_{11} - \gamma) \equiv 0, \quad y(\delta - \gamma) \equiv 0,$$

$$z(\beta - \alpha) \equiv 0, \quad w(b_{22} - \beta) \equiv 0,$$

where $x \not\equiv 0$, $w \not\equiv 0$. Hence when $\alpha \equiv \beta$, $\delta \equiv \gamma$ there results $a_{11} \equiv \gamma$, $b_{22} \equiv \alpha$. We are thus led to a single set of defining relations:

$$P_1^p = P_2^p = Q^q = R^r = 1, \quad P_1P_2 = P_2P_1, \quad Q^{-1}P_1Q = P_1^q,$$

$$Q^{-1}P_2Q = P_2^{q^k}, \quad R^{-1}P_1R = P_1^a, \quad R^{-1}P_2R = P_2^{a^h}, \quad RQ = QR,$$

$$\alpha^r \equiv 1(p), \quad \gamma^q \equiv 1(p) \quad (h = 1, 2, \dots, r-1; k = 1, 2, \dots, q-1),$$

or, briefly, say $G = (1 : \gamma 0 : 0\gamma^k : \alpha 0 : 0\alpha^h : 1)$. Proceeding to the determination of τ we observe that there are, by hypothesis, two subgroups, $\{P_1\}$, $\{P_2\}$, both permutable with Q and R . In any isomorphism of G with itself either $\{P_1\} \sim \{P_2\}$, $\{P_2\} \sim \{P_1\}$ or else $\{P_1\} \sim \{P_1\}$, $\{P_2\} \sim \{P_2\}$. Hence there are two choices of generators of order p . Every element of G is of the form $\Omega = R^x Q^y P_1^u P_2^v$. Hence $\Omega^s = R^{sx} Q^{sy} P_1^{us} P_2^{vs}$, so that Ω is of order r only when $y \equiv 0 \pmod{q}$ and of order q when $x \equiv 0 \pmod{r}$. Thus the most general operator of order q is $Q'_0 = Q^y P_1^u P_2^v$, which transforms G in the same manner as $Q_0 = Q^y$. Similarly $R'_0 = R^x$. Employing the new generators R_0, Q_0 , $P_{10} = P_1, P_{20} = P_2$, we get

$$(1 : \gamma 0 : 0\gamma^k : \alpha 0 : 0\alpha^h : 1) \sim (1 : \gamma^y 0 : 0\gamma^{ky} : \alpha^x 0 : 0\alpha^{hx} : 1).$$

Hence any set of relations involving arbitrary primitive roots (α^a, γ^b) can be transformed into the original set. Next let $P_{10} = P_2, P_{20} = P_1$. Then

$$(1 : \gamma 0 : 0\gamma^k : \alpha 0 : 0\alpha^h : 1) \sim (1 : \gamma^{ky} 0 : 0\gamma^{ky} : \alpha^{hx} 0 : 0\alpha^x : 1)$$

if

$$(6) \quad ky \equiv 1 \pmod{q}, \quad hx \equiv 1 \pmod{r}.$$

The group characterized by $[h, k]$ is thus isomorphic with $[x, y]$ when (6) is satisfied. Further τ equals the number of distinct solutions of (6), e. g., when $r = 2$, $\tau = \frac{1}{2}(q + 1)$, and when r is odd, $\tau = \frac{1}{4}(qr + q + r + 1)$.

[3] $i = \text{VI}$, $j = \text{V}$. When $h \neq 1$ we have $Q^{-1}P_jQ = P_j^{\alpha_j}$ ($j = 1, 2$). Assuming that

$$R^{-1}P'_1R = P_1^x P_2^y, \quad R^{-1}P'_2R = P_1^z P_2^w,$$

we derive

$$a_{11}x - z \equiv 0, \quad x - (\iota^p + \iota - a_{11})z \equiv 0,$$

$$a_{22}y - w \equiv 0, \quad y - (\iota^p + \iota - a_{22})w \equiv 0.$$

The elimination of x, y, z, w gives

$$a_{jj}^2 - (\iota^p + \iota)a_{jj} + 1 \equiv 0 \quad (j = 1, 2),$$

whence $a_{jj} = \iota^p$ or ι . Hence a_{11}, a_{22} are galoisian imaginaries* and G , for $i = \text{VI}$, $j = \text{V}$, does not exist.

Before considering the ambiguous case $h = 1$ a few general results must be established.

Let S and T be any set of generators of I_{p^2} , so that $G = \{S, T, Q, R\}$. We may write

$$P'_1 = S^x T^y, \quad P'_2 = S^z T^w,$$

$$Q^{-1}SQ = S^{a_{11}} T^{a_{21}}, \quad Q^{-1}TQ = S^{a_{12}} T^{a_{22}}.$$

Hence

$$Q^{-1}P'_1Q = P_2 = S^z T^w = S^{a_{11}x + a_{12}y} T^{a_{21}x + a_{22}y},$$

$$Q^{-1}P'_2Q = P_1^{-1}P_2^{\iota^p + \iota} = S^{-x + (\iota^p + \iota)z} T^{-y + (\iota^p + \iota)w} = S^{a_{11}z + a_{12}w} T^{a_{21}z + a_{22}w},$$

whence results the eliminant

$$D = \begin{vmatrix} x & y & z & w \\ a_{11} & a_{12} & -1 & 0 \\ a_{21} & a_{22} & 0 & -1 \\ 1 & 0 & a_{11} - t & a_{12} \\ 0 & 1 & a_{21} & a_{22} - t \end{vmatrix} \equiv 0 \pmod{p},$$

where $t = \iota^p + \iota$. Its expansion gives

$$D_{12}^2 - t(a_{11} + a_{22} - t)D_{12} + a_{22}^2 - a_{11}^2 + t(a_{11} - a_{22}) + 2a_{12}a_{21} + 1 \equiv 0.$$

Now assume $S = P_1$. Then, since $p \equiv -1 \pmod{q}$, $\rho_{q,p} = 0$ and we may take $Q^{-1}P_1Q \equiv U$ as T . Then

*SERRET, *Cours d'Algebre Supérieur*, cinq. ed. (1885), tome 2, sec. 3, chap. 3. See also DICKSON, *Linear Groups*, pp. 14-19.

$$J_Q = \left(\begin{array}{c} P_1^{x_1} U^{x_2} \\ P_1^{\alpha_{12} x_2} U^{x_1 + \alpha_{22} x_2} \end{array} \right), \quad J_Q^q = 1,$$

$$D_{12}^q = \begin{vmatrix} 0 & a_{12} \\ 1 & a_{22} \end{vmatrix}^q \equiv (-a_{12})^q \equiv 1 \pmod{p}.$$

Now $-a_{12}$ cannot be a primitive root of this congruence; for, if so $p \equiv 1 \pmod{q}$, whereas $p \equiv -1 \pmod{q}$ and $q > r$. It follows that $a_{12} \equiv -1 \pmod{p}$ and

$$D \equiv (a_{22} - \iota)^2 \equiv 0, \quad a_{22} \equiv \iota \equiv \iota^p + \iota.$$

This gives $I_{p^2} = \{P_1 U\}$ and

$$(7) \quad \begin{aligned} Q^{-1} P_1 Q &= U, & Q^{-1} U Q &= P_1^{-1} U^{\iota^p + \iota}, \\ R^{-1} P_1 R &= P_1^a, & R^{-1} U R &= P_1^\xi U^\eta, \end{aligned}$$

$$\delta_{12} = \begin{vmatrix} -1 & \xi \\ -\iota^p - \iota & \alpha - \eta \end{vmatrix} \equiv 0, \quad \delta'_{12} = \begin{vmatrix} 1 & 0 \\ -\iota^p - \iota & \alpha - \eta \end{vmatrix} \equiv 0,$$

and thus, when $h = 1$, $\eta \equiv \alpha$, $\xi \equiv 0 \pmod{p}$.

Inversely let $P_2 = P_1^{\xi'} U^{\eta'}$. Then

$$R^{-1} P_2 R = P_1^{\xi' a^h} U^{\eta' a^h} = P_1^{\xi' a} U^{\eta' a}$$

and hence $h = 1$. Thus when $h = 1$ there exists a group

$$G = \{P_1, U, Q, R\} = (1:01:-1\iota^p + \iota:\alpha 0:0\alpha:1),$$

where $\alpha^r \equiv 1 \pmod{p}$, $p \equiv 1 \pmod{r}$, $\tau = 1$. Also $p \equiv -1 \pmod{q}$ and, in the $GF[p^2]$, $\iota^q \equiv 1 \pmod{p}$.

[4] $i = V, j = VI$. Since r is necessarily an odd prime, the argument of [3] again gives for G a single type, $G = (1:\gamma 0:0\gamma:01:-1\iota^p + \iota:1)$, with $\gamma^q \equiv 1 \pmod{p}$, $p \equiv 1 \pmod{q}$, $\tau = 1$. Likewise $p \equiv -1 \pmod{r}$; and $\iota^r \equiv 1 \pmod{p}$ in the $GF[p^2]$.

[5] $i = VI, j = VI$. Employing as in [3] the theory of the determinant D we are led to the same equations (7), viz.,

$$Q^{-1} P_1 Q = U, \quad Q^{-1} U Q = P_1^{-1} U^{\iota_1^p + \iota_1}, \quad \iota_1^q \equiv 1 \pmod{p}.$$

Let us assume that

$$R^{-1} P_1 R = P_2 = P_1^x U^y, \quad R^{-1} U R = P_1^z U^w.$$

Then

$$\delta_{12} = \begin{vmatrix} -1 & z \\ -\iota_1^p - \iota_1 & x - w \end{vmatrix} \equiv 0, \quad \delta'_{12} = \begin{vmatrix} 1 & y \\ -\iota_1^p - \iota_1 & x - w \end{vmatrix} \equiv 0,$$

$$d_{12} = \begin{vmatrix} -1 & 1 \\ z & y \end{vmatrix} \equiv 0, \quad D_{12} = \begin{vmatrix} x & z \\ y & w \end{vmatrix} \not\equiv 0.$$

Thus

$$z \equiv -y, \quad w \equiv x + (\iota_1^p + \iota_1)y, \quad D_{12} \equiv x^2 + (\iota_1^p + \iota_1)xy + y^2.$$

Since

$$R^{-1}P_2R = P_1^{-1}P_2^{\iota_2^p + \iota_2}, \quad \iota_2^r \equiv 1(p),$$

so that

$$R^{-1}UR = P_1^{-y^2}Uxy + (\iota_1^p + \iota_1)y^2 = P_1^{-(x^2+1) + (\iota_2^p + \iota_2)x}U^{-xy + (\iota_2^p + \iota_2)y}.$$

Since P_1 and P_2 are independent, $y \not\equiv 0$; hence

$$(8) \quad 2x + (\iota_1^p + \iota_1)y - (\iota_2^p + \iota_2) \equiv 0,$$

$$(9) \quad y^2 - x^2 + (\iota_2^p + \iota_2)x - 1 \equiv 0.$$

From the latter we at once derive

$$D_{12} = x^2 + (\iota_1^p + \iota_1)xy + y^2 \equiv 1,$$

$$(10) \quad (\iota_2 - \iota_1^2 \iota_2)^2 x^2 - (1 - \iota_1^2)(\iota_2 - \iota_2^3)x + (1 - \iota_1^2 \iota_2^2)(\iota_2^2 - \iota_1^2) \equiv 0,$$

$$(11) \quad (\iota_1 - \iota_1^p)^2 y^2 - (\iota_2 - \iota_2^p)^2 \equiv 0.$$

There always exist integral solutions of (10) and (11), $x = \epsilon_j, y = \sigma_j (j = 1, 2)$.

Thus

$$R^{-1}P_1R = P_1^{\epsilon_j + (\iota_1^p + \iota_1)\sigma_j}U^{-\sigma_j}, \quad R^{-1}UR = P_1^{\sigma_j}U^{\epsilon_j}.$$

THEOREM. *The two general types of G characterized by the two distinct sets of solutions of (10) and (11), viz. $[\epsilon_1, \sigma_1]$ and $[\epsilon_2, \sigma_2]$ are simply isomorphic.*

In proof, $\sigma_2 \equiv -\sigma_1$, and congruence (8) gives

$$2\epsilon_2 - (\iota_1^p + \iota_1)\sigma_1 - (\iota_2^p + \iota_2) \equiv 0, \quad \epsilon_2 \equiv \epsilon_1 + (\iota_1^p + \iota_1)\sigma_1.$$

Hence the two types of G are characterized by

$$R^{-1}P_1R = P_1^{\epsilon_1 + (\iota_1^p + \iota_1)\sigma_1}U^{-\sigma_1}, \quad R^{-1}UR = P_1^{\sigma_1}U^{\epsilon_1},$$

and

$$R^{-1}P_1R = P_1^{\epsilon_1}U^{\sigma_1}, \quad R^{-1}UR = P_1^{-\sigma_1}U^{\epsilon_1 + (\iota_1^p + \iota_1)\sigma_1}.$$

Let us select a new operation of order q from $\{Q\}$, e. g. $Q' = Q^{-1}$. Then $Q'R = RQ'$, $Q'^{-1}UQ' = P_1$,

$$Q'^{-1}P_1Q' = U^{r_1}P_1^{r_2} = U^{-1}P_1^{\iota_1^p + \iota_1}, \quad r_j = \frac{\iota_1^{(q-j)p} - \iota_1^{q-j}}{\iota_1^p - \iota_1}.$$

The result of selecting Q' and (ϵ_2, σ_2) is thus to interchange P_1 and U and to reproduce the relations given by Q and (ϵ_1, σ_1) . Hence $[\epsilon_2, \sigma_2] \sim [\epsilon_1, \sigma_1]$.

The quantities ι_1 and ι_2 are marks of the $GF[p^2]$ and in that field appertain

respectively to the exponents q and r . Let ρ be any primitive root in the $GF[p^2]$. It is easy to show that $\tau = 1$ and hence we may select *

$$\iota_1 \equiv \rho^{(p^2-1)/q}, \quad \iota_2 \equiv \rho^{(p^2-1)/r},$$

thus

$$G = (1 : 01 : -1, \iota_1^p + \iota_1 : \epsilon + (\iota_1^p + \iota_1)\sigma, -\sigma : \sigma\epsilon : 1),$$

where

$$\iota_1 \equiv \rho^{(p^2-1)/q}, \quad \iota_2 \equiv \rho^{(p^2-1)/r}, \quad \rho^{p^2-1} \equiv 1; \quad p \equiv -1 \pmod{qr}, \quad \tau = 1,$$

$$(\iota_1 - \iota_1^p)^2 \sigma^2 - (\iota_2 - \iota_2^p)^2 \equiv 0, \quad 2\epsilon + (\iota_1^p + \iota_1)\sigma - (\iota_2^p + \iota_2) \equiv 0.$$

§ 3. The generating function $[k]$.

Consider the relation $R^{-z}P_1R^z = P_1^{u_z}U^{v_z}$. From it

$$u_{z+1} - (2x + t_1y)u_z + (x^2 + t_1xy + y^2)u_{z-1} \equiv 0,$$

$$u_{z+1} - t_2u_z + u_{z-1} \equiv 0 \quad (t_j = \iota_j^p + \iota_j; j=1, 2),$$

These recurring formulæ give

$$u_k \equiv [k]_2x - [k-1]_2, \quad v_k \equiv [k]_2y,$$

where

$$[k]_j \equiv \frac{\iota_j^{kp} - \iota_j^k}{\iota_j^p - \iota_j}.$$

Following are some of the properties of the generating function $[k]_j$.

$$(12) \quad \frac{[k+1]_j}{[k]_j} = \frac{1}{t_j} + \frac{1}{t_j} + \frac{1}{t_j} + \dots k \text{ terms},$$

$$(13) \quad [k]_j^2 - [k+1]_j[k-1]_j - 1 \equiv 0,$$

$$(14) \quad [0]_j \equiv 0, \quad [1]_j \equiv 1, \quad [-k]_j \equiv -[k]_j,$$

$$(15) \quad [k+1]_j \equiv [2]_j[k]_j - [k-1]_j,$$

$$(16) \quad \{[k+1]_j - [k-1]_j - [2]_j\} \iota_j^k \equiv (\iota_j^{k+1} - 1)(\iota_j^{k-1} - 1).$$

§ 4. Class (10), $p > q > r$.

We shall consider next groups possessing a single maximal self-conjugate subgroup $H_{p^2q, i}$ of non-abelian type ($i = \text{III, IV, V, VI}$). It is readily shown that class (10, 12), with $i = \text{III}$, must contain an invariant subgroup H_{p^2r} . Class (10) remains to be considered.

[1] $i = \text{IV}$. Here $H_{p^2q, \text{IV}} = \{P, Q\}$ and since $\{P\}$ is self-conjugate in G , $R^{-1}PR = P^\beta$. Since $\rho_{R, q} \geq 1$ [Eq. (1)], $R^{-1}QR = Q^\gamma$. Hence

$$(QR)^{-1}P(QR) = P^{\alpha\beta} = (RQ^\gamma)^{-1}P(RQ^\gamma) = P^{\beta\alpha\gamma}, \quad \alpha^q \equiv 1 \pmod{p^2},$$

$$\alpha\beta(\alpha^{\gamma-1} - 1) \equiv 0 \pmod{p^2}, \quad \gamma \equiv 1 \pmod{q}.$$

* DICKSON, *Linear Groups*, p. 13.

Hence $\{P_1, P_2, R\}$ is self-conjugate in $\{P_1, P_2, Q, R\} = G$, contrary to hypothesis.

[2] $i = V$. Let $H_{p^2q, V} = \{P'_1, P_2, Q\}$. Assuming that

$$R^{-1}P'_1R = P_1^{a_{11}}P_2^{a_{21}}, \quad R^{-1}P_2R = P_1^{a_{12}}P_2^{a_{22}},$$

we deduce

$$\alpha_{11}\alpha(\alpha^{\gamma-1} - 1) \equiv 0, \quad a_{21}(\beta^{\gamma} - \alpha) \equiv 0,$$

$$a_{22}\beta(\beta^{\gamma-1} - 1) \equiv 0, \quad a_{12}(\alpha^{\gamma} - \beta) \equiv 0,$$

where $\alpha^q \equiv 1(p)$, $\beta \equiv \alpha^h$. Now $\gamma \not\equiv 1 \pmod{q}$. Hence

$$\alpha_{11} \equiv 0, \quad a_{22} \equiv 0, \quad \alpha^{\gamma h} \equiv \alpha, \quad \alpha^{\gamma} \equiv \alpha^h \pmod{p},$$

$$\gamma \equiv h \pmod{q}, \quad \alpha^{\gamma^2} \equiv \alpha \pmod{p}, \quad \gamma^2 \equiv 1 \pmod{q}.$$

But γ appertains to the exponent r modulo q , and therefore $r = 2$ and $\gamma \equiv -1 \pmod{q}$. Thus

$$R^{-1}P'_1R = P_2^{a_{21}}, \quad R^{-1}P_2R = P_1^{a_{12}}, \quad a_{12}a_{21} \equiv 1 \pmod{p}.$$

Then $P_1 = P_1^{a_{12}}$, P_2, Q, R , generate a group of order $2p^2q$, viz., $G = (1 : \alpha 0 : 0 \alpha^{\gamma-1} : 0 1 : 10 : -1)$. Also $p \equiv 1(q)$, $\tau = 1$.

[3] $i = VI$. It has been shown [§ 1], that $p \equiv \pm 1 \pmod{r}$.

(a) First let $p \equiv 1(r)$. Then $\rho_{R,p} \cong 2$ and two subgroups $\{P_1\}, \{P_3\}$ may be selected which are permutable with R . If

$$Q^{-1}P_1Q = P_2, \quad Q^{-1}P_2Q = P_1^{-1}P_2^{p+1},$$

then

$$R^{-1}P_1R \equiv P_1^{\beta}, \quad R^{-1}QR = Q^{\gamma}, \quad \gamma \not\equiv 1 \pmod{q}.$$

Since I_{p^2} is invariant in G we may assume that

$$P_3 = P_1^zP_2^v, \quad R^{-1}P_2R = P_1^xP_2^y,$$

Hence

$$(QR)^{-1}P_1(QR) = P_1^xP_2^y = (RQ^{\gamma})^{-1}R_1(RQ^{\gamma}) = P_1^{-\beta[\gamma-1]}P_2^{\beta[\gamma]},$$

$$(QR)^{-1}P_2(QR) = P_1^{-\beta+[2]x}P_2^{[2]y} = (RQ^{\gamma})^{-1}P_2(RQ^{\gamma}) = P_1^{-[\gamma-1]x-[\gamma]y}P_2^{[\gamma]x+[\gamma+1]y},$$

$$x \equiv -[\gamma-1]\beta, \quad y \equiv [\gamma]\beta,$$

$$[\gamma]^2 \equiv [\gamma-1]^2 + [2][\gamma-1] + 1,$$

$$[\gamma]\{[\gamma+1] - [\gamma-1] - [2]\} \equiv 0.$$

Now $[\gamma] \not\equiv 0 \pmod{q}$. Since $[-k] \equiv -[k]$ and

$$[\gamma+1] - [\gamma-1] - [2] \equiv (\nu^{\gamma+1} - 1)(\nu^{\gamma-1} - 1) \equiv 0 \text{ [Eq. (16)]},$$

there results $\gamma \equiv -1 \pmod{q}$, $\gamma^r \equiv (-1)^r \equiv +1 \pmod{q}$, whence $r = 2$. If $R^{-1}P_3R = P_3^a$, then $\alpha \equiv \pm 1 \pmod{p}$,

$$\begin{aligned} w(y \mp 1) &\equiv 0, & xw + z(\beta \mp 1) &\equiv 0, \\ w(-\beta \mp 1) &\equiv 0, & [2]\beta w + z(\beta \mp 1) &\equiv 0. \end{aligned}$$

First let the upper sign hold. If $\beta \equiv 1$, then $w \equiv 0$ which is impossible, since P_1, P_3 are independent. Hence $\beta \equiv -1$, $x \equiv -[2]$, $y \equiv +[1] \equiv +1$. Likewise if we use the lower sign, $\beta \equiv +1$, $x \equiv +[2]$, $y \equiv -[1] \equiv -1$. We thus obtain the two sets of defining relations:

$$(1:01:-1\iota^p + \iota:\mp 10:\iota^{\mp p} + \iota^{\mp 1}, \pm 1:-1).$$

To determine τ , let $Q_0 = Q^x$, $R_0 = R$, $P_{1_0} = P_1$, $P_{2_0} = P_1^{[x-1]}P_2^{[x]}$; there results

$$\{P_{1_0}, P_{2_0}, Q_0, R_0\} = (1:01:-1\iota^{xp} + \iota^x:\mp 10:\pm[x-1]\mp[2][x], \pm[x]:-1).$$

But

$$\pm[x-1]\mp[2][x] \equiv \mp[x+1] \equiv \mp(\iota^{xp} + \iota^x)\mp[x-1],$$

[Eq. (15)]. Hence

$$\{P_{1_0}, P_{2_0}, Q_0, R_0\} = (1:01:-1\iota^{xp} + \iota^x:\mp 10:\mp(\iota^{xp} + \iota^x), \pm 1:-1) \sim G.$$

Thus the same defining relations are reproduced with ι replaced by ι^x , and so $\tau = 1$.

It will now be proved that these two types are simply isomorphic. Select new operators as follows:

$$q_1 = Q, r_1 = R, p_1 = P_1^a P_2^b, p_2 = P_1^{-b} P_2^{a+[2]b} = q_1^{-1} p_1 q_1.$$

Then using the first set of defining relations we will have

$$q_1^{-1} p_2 q_1 = p_1^{-1} p_2^{\iota^p + \iota}, r_1^{-1} p_1 r_1 = p_1, r_1^{-1} p_2 r_1 = p_1^{\iota^p + \iota} p_2^{-1}, r_1^{-1} q_1 r_1 = q_1^{-1}$$

if

$$2a + [2]b \equiv 0 \pmod{p}.$$

Hence when a new operator $p_1 = P_1^a P_2^b$ is selected, where a and b are solutions of $2a + (\iota^p + \iota)b \equiv 0 \pmod{p}$, the first type is transformed into the second. They are therefore isomorphic.

(b) When $p \equiv -1(r)$, r odd, $\rho_{R,p} = 0$. As before, we deduce

$$\begin{aligned} Q^{-1}P_1Q &= P_2, & Q^{-1}P_2Q &= P_1^{-1}P_{2_-}^{\iota_1^p + \iota_1}, & \iota_1^q &\equiv 1(p), \\ R^{-1}P_1R &= P_3, & R^{-1}P_3R &= P_1^{-1}P_{3_-}^{\iota_2^p + \iota_2}, & \iota_2^r &\equiv 1(p), \end{aligned}$$

Let $P_3 = P_1^x P_2^y$ and $R^{-1}P_2R = P_4 = P_1^z P_2^w$. Then

$$(17) \quad R^{-1}P_2R = P_1^{-(x^2+1)+[2]_2x} P_2^{y[2]_2-yx} = P_1^{[\gamma-1]_1xy-[y]_1y^2} P_2^{[\gamma]_1xy+[y+1]_1y^2}.$$

In addition to the latter, but not independent of them, we have the congruences derived from

$$(18) \quad (QR)^{-1}P_2^y(QR) = (RQ^y)^{-1}P_2^y(RQ^y).$$

The equations (17) and (18) give us the dialytic eliminant

$$\Delta_{12} = \{\iota_2^p + \iota_2\} \{[\gamma]_1^2 - (\iota_2^{2p} + \iota_2^2)[\gamma]_1 + 1\} \{(\iota_1^{r+1} - 1)(\iota_1^{r-1} - 1)\}^2 \equiv 0.$$

Now $[\gamma]_1$ is an integer, and since $r \neq 2$, and $\gamma \neq -1$, it follows that $\gamma \equiv 1 \pmod{q}$, contrary to hypothesis. Hence when $p \equiv -1 \pmod{r}$ and r is odd, no corresponding group G exists.

The results of this section may be summarized in the following

THEOREM. *A group G_{p^2qr} ($p > q > r$) always contains a maximal self-conjugate subgroup H of order p^2q . If H is the only maximal invariant subgroup of G and if r is odd, then $N_q = 1$ and H is necessarily abelian. If r is even ($r = 2$) and $p \equiv 1 \pmod{q}$ there exists one type whose subgroup H_{p^2q} is non-abelian, and if r is even and $p \equiv -1 \pmod{q}$ there exists a second type possessing a non-abelian H_{p^2q} . These two types of G contain respectively q and pq operators (and subgroups) of order 2, and in each type $N_q = p^2$. Moreover, with exception of the two types just described, every group of order p^2qr ($p > q > r$), in which $N_r \equiv 0 \pmod{q}$, possesses an abelian maximal self-conjugate subgroup H_{p^2q} .*

A general summary of all the existent types of G follows. Except for ι and ρ , every parameter occurring in the tables is an integer; while ι and ρ are marks of the $GF[p^2]$. See footnote on the second page of the paper.

TABLE 1. $p > q > r$.

I_p non-cyclical; $P_1^n = P_2^n = Q^r = R^r = 1$, $P_1 P_2 = P_2 P_1$. I_p cyclical; $P_1^p = Q^r = R^r = 1$.
Case (a), $QR = RQ$.

Class	$Q^{-1}P_1Q$	$Q^{-1}P_2Q$	$R^{-1}P_1R$	$R^{-1}P_2R$	Parameters.	Arith. Rel.	τ
[12...12]	P_1	P_1	P_1	P_1	\cdot	\cdot	1
"	P_1	P_2	P_1	P_2	\cdot	\cdot	1
[3467891012]	P_1^a	P_2	P_1	P_2	$\alpha^q \equiv 1(p)$	$p \equiv 1(q)$	1
[3489101112]	P_1	P_2	P_1^a	P_2	$\alpha^r \equiv 1(p)$	$p \equiv 1(r)$	1
[46891012]	P_1^a	P_2	P_1^{β}	P_2	$\alpha^q \equiv \beta^r \equiv 1(p)$	$p \equiv 1(qr)$	1
[78910]	P_1^a	\cdot	P_1	\cdot	$\alpha^q \equiv 1(p^2)$	$p \equiv 1(q)$	1
"	P_1^a	$P_2^{a^h}$	P_1	P_2	$\alpha^q \equiv 1(p)$	$p \equiv 1(q)$	$\frac{1}{2}(q+1)$
"	P_2	$P_1^{-1}P_2^{p+1}$	P_1	P_2	$\ell^q \equiv 1(p)$	$p \equiv -1(q)$	1
[891012]	P_1^a	P_2	P_1	P_2^{β}	$\alpha^q \equiv \beta^r \equiv 1(p)$	$p \equiv 1(qr)$	1
[9101112]	P_1	\cdot	P_1^a	\cdot	$\alpha^r \equiv 1(p^2)$	$p \equiv 1(r)$	1
"	P_1	P_2	P_1^a	$P_2^{a^h}$	$\alpha^r \equiv 1(p)$	$p \equiv 1(r)$	1 or $\frac{1}{2}(r+1)$
"	P_1	P_2	P_2	$P_1^{-1}P_2^{p+1}$	$\ell^r \equiv 1(p)$	$p \equiv -1(r)$	1
[8910]	P_1^{β}	$P_2^{a^h}$	P_1^a	P_2	$\alpha^r \equiv \beta^q \equiv 1(p)$	$p \equiv 1(qr)$	$q-1$
[91012]	P_1^a	P_2	P_1^{β}	$P_2^{a^h}$	$\alpha^q \equiv \beta^r \equiv 1(p)$	$p \equiv 1(qr)$	$r-1$
[910]	P_1^a	\cdot	P_1^{β}	\cdot	$\alpha^q \equiv \beta^r \equiv 1(p^2)$	$p \equiv 1(qr)$	1
"	P_1^{γ}	$P_2^{\gamma^k}$	P_1^a	$P_2^{a^h}$	$\gamma^q \equiv \alpha^r \equiv 1(p)$	$p \equiv 1(qr)$	$\frac{1}{2}(q+1)$ or $\frac{1}{4}(r+1)(q+1)$
"	P_2	$P_1^{-1}P_2^{p+1}$	P_1^a	P_2^a	$\ell^q \equiv 1(p)$	$p \equiv -1(q)$	1
"					$\alpha^r \equiv 1(p)$	$p \equiv 1(r)$	
"	P_2	$P_1^{-1}P_2^{p^2-1}$	P_1^a	P_2^a	$\rho = \text{prim. root in } GF[p^2]; \iota_1, \iota_2 = \rho^{(p^2-1)/q, r}$	$p \equiv -1(qr)$	1
"	P_2	$P_1^{-1}P_2^{\frac{p^2-p}{q} + p}$	P_1^a	$P_2^{-\sigma}$	$2\epsilon + [2]_1 \sigma - [2]_2 \equiv 0$		
"	P_2	$P_1^{-1}P_2^{\frac{p^2-p}{q} + p}$	P_1^a	$P_2^{-\sigma}$	$(\iota_1 - \iota_1^r)^2 \sigma^2 - (\iota_2 - \iota_2^r)^2 \equiv 0$		
"	P_1^{γ}	P_2^{γ}	P_2	$P_1^{-1}P_2^{p+1}$	$\ell^r \equiv 1(p)$	$p \equiv -1(r)$	1
"					$\gamma^q \equiv 1(p)$	$p \equiv 1(q)$	

Case (b). $R^{-1}QR = Q^h$; $\gamma^r \equiv 1(q)$.

Class.	$Q^{-1}P_1Q$	$Q^{-1}P_2Q$	$R^{-1}P_1R$	$R^{-1}P_2R$	Parameters.	Arith. rel.	τ
[256101112]	P_1	P_2	P_1	P_2	$h = 1$	$q \equiv 1(r)$	1
"	P_1	.	P_1	.	$h = 1$	$q \equiv 1(r)$	1
[56101112]	P_1	P_2	P_1^a	P_2	$h = 1, 2 \dots r-1$ $\alpha^r \equiv 1(p)$	$p \equiv q \equiv 1(r)$	$r-1$
[101112]	P_1	.	P_1^a	.	$h = 1, 2 \dots r-1$ $\alpha^r \equiv 1(p^2)$	$p \equiv q \equiv 1(r)$	$r-1$
"	P_1	P_2	P_1^a	P_2^k	$h, k=1, 2 \dots r-1$ $\alpha^r \equiv 1(p)$	$p \equiv q \equiv 1(r)$	1 or $\frac{1}{2}(r^2-1)$
"	P_1	P_2	P_2	$P_1^{-1}P_2^{p+1}$	$h = 1, 2 \dots r-1$ $\iota^r \equiv 1(p)$	$p \equiv -q \equiv -1(r)$	$r-1$
[10]	P_1^a	$P_2^{a^{q-1}}$	P_2	P_1	$h = 1, \gamma \equiv -1$ $\alpha^q \equiv 1(p)$	$r = 2$ $p \equiv 1(q)$	1
"	P_2	$P_1^{-1}P_2^{p+1}$	P_1^{-1}	$P_1^{-p+1}P_2$	$h = 1, \gamma \equiv -1$ $\iota^q \equiv 1(p)$	$r = 2$ $p \equiv -1(q)$	1

TABLE 2. $q > p > r$.

I_{p^2} non-cyclical; $P_i^p = Q^q = R^r = 1$ ($i = 1, 2$), $P_1P_2 = P_2P_1$, $RP_2 = P_2R$,
 I_{p^2} cyclical; $P_1^{p^2} = Q^q = R^r = 1$, $RP_1 = P_1R$.

Class.	$P_1^{-1}QP_1$	$P_2^{-1}QP_2$	$R^{-1}QR$	$R^{-1}P_1R$	Parameters.	Arith. Rel.	τ
[1234561112]	Q^a	.	Q	P_1	$\alpha^p \equiv 1(q)$	$q \equiv 1(p)$	1
[12511]	Q^a	.	Q	P_1	$\alpha^{p^2} \equiv 1(q)$	$q \equiv 1(p^2)$	1
[1112]	Q^a	.	Q^r	P_1	$\alpha^p \equiv \gamma^r \equiv 1(q)$	$q \equiv 1(pr)$	1
[11]	Q^a	.	Q^r	P_1	$\alpha^{p^2} \equiv \gamma^r \equiv 1(q)$	$q \equiv 1(p^2r)$	1
[1251112]	Q	Q^a	Q	P_1	$\alpha^p \equiv 1(q)$	$q \equiv 1(p)$	1
[251112]	Q	Q^r	Q^a	P_1	$\gamma^p \equiv \alpha^r \equiv 1(q)$	$q \equiv 1(pr)$	1
[4561112]	Q	Q^a	Q	P_1^δ	$\alpha^p \equiv 1(q)$ $\delta^r \equiv 1(p)$	$q \equiv 1(p)$ $p \equiv 1(r)$	1
[561112]	Q	Q^a	Q^{r^h}	P_1^δ	$\alpha^p \equiv \gamma^r \equiv 1(q)$ $\delta^r \equiv 1(p)$	$q \equiv 1(pr)$ $p \equiv 1(r)$	$r-1$

TABLE 3. $q > r > p$.*Case (a).*

I_{p^2} non-cyclical; $P_i^p = Q^q = R^r = 1 (i = 1, 2)$, $P_1P_2 = P_2P_1$, $RQ = QR$,
 I_{p^2} cyclical; $P_1^{p^2} = Q^q = R^r = 1$, $QR = RQ$.

Class.	$P_1^{-1}QP_1$	$P_2^{-1}QP_2$	$P_1^{-1}RP_1$	$P_2^{-1}RP_2$	Parameters.	Arith. Rel.	τ
[12345678]	Q	.	R^a	.	$\alpha^p \equiv 1(r)$	$r \equiv 1(p)$	1
[1237]	Q	.	R^a	.	$\alpha^{p^2} \equiv 1(r)$	$r \equiv 1(p^2)$	1
[123456]	Q^a	.	R^{β^h}	.	$\alpha^p \equiv 1(q)$ $\beta^p \equiv 1(r)$	$q \equiv r \equiv 1(p)$	$p - 1$
[125]	Q^a	.	R^{β^h}	.	$\alpha^{p^2} \equiv 1(q)$ $\beta^p \equiv 1(r)$	$q \equiv 1(p^2)$ $r \equiv 1(p)$	$p - 1$
[234]	Q^{a^h}	.	R^β	.	$\alpha^p \equiv 1(q)$ $\beta^{p^2} \equiv 1(r)$	$r \equiv 1(p^2)$ $q \equiv 1(p)$	$p - 1$
[12]	Q^a	.	R^{β^h}	.	$\alpha^{p^2} \equiv 1(q)$ $\beta^{p^2} \equiv 1(r)$	$q \equiv r \equiv 1(p^2)$	$p^2 - 1$
[12345678]	Q	Q	R	R^a	$\alpha^p \equiv 1(r)$	$r \equiv 1(p)$	1
[123]	Q	Q^a	R	R^{β^h}	$\alpha^p \equiv 1(q)$ $\beta^p \equiv 1(r)$	$q \equiv r \equiv 1(p)$	$p - 1$
[1235]	Q	Q^β	R^a	R	$\alpha^p \equiv 1(r)$ $\beta^p \equiv 1(q)$	$q \equiv r \equiv 1(p)$	1

Case (b). The simple group $G_{15!}$, $p = 2$, $q = 5$, $r = 3$.

$$Q^5 = 1, \quad P^2 = 1, \quad (QP)^3 = 1, \quad [R = QP].$$